

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/000833

International filing date: 26 January 2005 (26.01.2005)

Document type: Certified copy of priority document

Document details: Country/Office: EP  
Number: 04075294.1  
Filing date: 30 January 2004 (30.01.2004)

Date of receipt at the International Bureau: 28 February 2005 (28.02.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr.    Patent application No.    Demande de brevet n°

04075294.1

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk





Anmeldung Nr:  
Application no.: 04075294.1  
Demande no:

Anmeldetag:  
Date of filing: 30.01.04  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

DSM IP Assets B.V.  
Het Overloon 1  
6411 TE Heerlen  
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Carboxypeptidase for cheese ripening

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

A23C/

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PT RO SE SI SK TR LI



EPO - DG 1

30. 01. 2004

**CARBOXYPEPTIDASE FOR CHEESE RIPENING**

(61)

**Field of the invention**

The present invention relates to cheese ripening.

**Background of the invention**

Flavour of food products is one of the key attributes for the consumer. In fermented products, e.g. dairy products, flavors are derived from milk components by enzymatic activities of micro-organisms. In cheese for instance, various flavour compounds have been identified as being essential and many of them are derived from casein degradation. Other enzymatic processes, such as lipolysis, are also involved, most notably in cheese where fungi are involved in the ripening process, e.g. Camembert and Roquefort cheese. In addition, lactose fermentation might lead to the flavour compounds such as propionic acid (Smit *et al*, Food Res. Int. (2000) **33**, 153-160).

Proteolysis in cheese during ripening plays a vital role in the development of texture as well as flavour and has been subject of several reviews (see e.g. McSweeney & Sousa, Lait (2000) **80**, 293-324). Proteolysis contributes to textural changes of the cheese matrix, due to breakdown of the protein network, decrease in  $a_w$ , through water binding by liberated carboxyl and amino groups and increase in pH, which facilitates the release of sapid compounds during mastication (Sousa et al, Int. Dairy Journal (2001), **11**, 327-345). It contributes directly to flavour and to off-flavour (e.g. bitterness) of cheese through the formation of peptides and free amino acids as well as liberation of substrates (amino acids) for secondary catabolic changes, i.e. transamination, deamination, decarboxylation, desulphuration, catabolism of aromatic amino acids and reactions of amino acids with other compounds. The rate and pattern of proteolysis may be influenced by location within the cheese.

Cheese ripening is a time-consuming process involving complex and well-balanced reactions between glycolysis, proteolysis and lipolysis of the milk components. In most cheeses, bacterial enzymes play a major role in this process. It is well known that changing the bacterial enzyme content is cheese directly affects the rate of cheese ripening and its final flavour (Klein & Lortal, Int. Dairy Journal (1999) **9**, 751-762.). A way

to influence cheese ripening is to increase the bacterial enzyme pool in cheese curd by the addition of whole lactic acid bacteria, unable to grow and produce significant levels of lactic acid, but still delivering active ripening enzymes during cheese ageing. The starters are normally weakened and referred to as attenuated.

5        Since cheese ripening is a time consuming process it is also costly. Cheeses need to be stored during ripening under precisely defined conditions for temperature and humidity for weeks to months. The ripening time varies considerably between the various cheeses, from 3 weeks (e.g. Mozzarella) to more than 2 years (e.g. Parmesan, extra mature cheddar). Any process that would result in acceleration of cheese ripening is  
10       interesting from an economic point of view: the same amount of cheese can be produced in a shorter time interval.

      Proteolysis in cheese is a very complex process, and proteases from various origins are involved (for review see e.g. Fox & McSweeney, Food Rev. Int (1996) 12, 457-509). Such proteases are the coagulant that was used during cheese manufacture  
15       (e.g. chymosin, pepsin or fungal acid proteinases), milk own proteins (e.g. plasmin), the proteases provided by the starter bacteria, proteases from non-starter adventitious microflora, proteases from a second inoculum (in some varieties, e.g. *P. roqueforti*, *P. camemberti*, *Br. Linens*), attenuated bacterial cells and exogenous proteases. The attenuated cells and the exogenous proteases are recent tools in the development of  
20       acceleration of cheese ripening. The generation of free amino acids is an important step in the acceleration of cheese ripening. Although the free amino acids contribute to the overall cheese flavour, their contribution is relatively small. The amino acids are the precursors, which are subsequently converted by the micro-organisms that are present in the cheese to flavour compounds. Availability of amino acids is therefore important for  
25       cheese flavour formation and thus for cheese ripening.

      The main alternative to the use of natural cheese in processed consumer foods requiring a cheese flavour are high-intensity cheese flavour concentrates such as enzyme-modified cheese (EMCs), cheese powders and cheese flavours (Kilcawley, Wilkinson & Fox, Int. Dairy Journal (1998), 8, 1-10). Cheese flavours are not produced  
30       for use in cheese as such but are produced to be applied in other food that conventionally contains natural cheese. Cheese is traditionally added to products as a spray dried preparation for flavour, appearance and texture enhancement. The amount of cheese used varies, and the overall flavour and quality of the product depends on the

type of cheese used. Cheese that has been treated enzymatically to enhance the flavour or a significant portion of the profile of that cheese is considered to be an EMC and it provides the food manufacturer with a strong cheese note in a form that is cost effective, nutritious and natural (Kilcawley, Wilkinson & Fox, *Int. Dairy Journal* (1998), **8**, 1-10).

5 EMCs have a flavour profile, which may be quite different from that of a natural cheese and yet on dilution with a suitable bland or nearly bland base, provide the desired cheesy note in the end product. The basis of EMC technology is the use of specific enzymes to produce typical cheese flavours from suitable substrates. Proteases, which term includes both endo-proteases and exo-proteases, are important enzymes in EMC production.  
10 Their role is similar to that in cheese ripening. Proteolysis in EMC is extensive and produces both high levels of savoury and bitter notes; the latter can be prevented, removed or masked by either controlled proteolysis, addition of specific exopeptidases or inclusion of masking agents, e.g. mature cheese or monosodium glutamate (Wilkinson & Kilcawley, *Bulletin of the IDF* (2002), **371**, 10-15).

15 Many proteases are involved in the generation of cheese flavours. The generation of the proper flavour for a cheese (or cheese derived product like EMC) requires a delicate balance of proteolytic activities from the proteases involved. Any imbalance will easily lead to flavours that are not wanted, such as bitterness development. Especially the development of bitterness in cheese (or EMC) has been  
20 well described and documented (see e.g. LeMieux & Simard, *Lait* (1991) **71**, 599-636; Lemieux & Simard, *Lait* (1992) **72**, 335-382) The development of proteases for cheese or EMC to enhance ripening processes is therefore a very delicate and complicated process. Exo-proteases are preferred over endo-proteases because they have a lower tendency to induce formation of bitterness. Endo-proteases are known to easily  
25 introduce such bitterness and are therefore preferably not used. There is, however, a clear industrial need for cheese ripening enzymes such as proteases, and over the years several commercial protease preparation have been introduced into the market. An overview of available commercial products is given in several papers (Wilkinson, van den Berg & Law, *Bulletin of the IDF* (2002) **371**, 16-19; Kilcawley, Wilkinson & Fox, *Food Biotechnol* (2002) **16**, 29-55; Kilcawley, Wilkinson & Fox, *Enzyme Microb. Technol.*  
30 (2002) **31**, 310-320). Examples include enzyme preparations derived from fungal species (including but not limited to Bioprotease P Conc and Bioprotease A conc from Quest, The Netherlands, Protease M & Protease A and Acid protease A from Amano, Promod



215 from Biocatalysts, Sternzyme B5026 from Stern, Flavourzyme MG/A from  
Novozymes, Denmark) and bacterial species (including but not limited to Protamex and  
Neutrase from Novozymes, Denmark, Protease N from Amano, Promod 24P and 24L  
from Biocatalysts, Protease B500 from DSM, The Netherlands, and Protease 200L from  
5 Rhodia Foods, France). The proteases vary considerably in composition with respect to  
presence of specific proteases and/or the ratio in which these proteases occur in a  
specific product. The papers by Kilcawley, Wilkinson and Fox, referred to above, clearly  
show that most commercial protease products are a mixture of endo- and exo-peptidase  
activities. Several products are developed to contain only exo-peptidase activity. For food  
10 applications, these are invariably amino-peptidases, and examples include DBS50 and  
DBP20 (from Rhodia, France), Corolase LAP (from Rohm, Germany), Flavourzyme  
MG/A (from Novozymes, Denmark), Accellerzyme AP (from DSM, The Netherlands) and  
Peptidase R (from Amano, Japan). The amino peptidases are developed and selected  
for the release amino acids that are important precursors of cheese flavour such as  
15 leucine, phenylalanine and valine. Several patent applications (e.g. WO96/38549)  
describe the preparation and use of amino peptidases, free from endo-proteases, which  
can be used to accelerate cheese ripening. Although there is a description of the use of a  
single wheat carboxy-peptidase to reduce the bitterness of bitter peptides from milk  
casein peptides (Umetsu, Matsuoka & Ichishima, J. Agric. Food Chem (1983)31, 50-53),  
20 there is no description of the use of a protease preparation containing a single carboxy-  
peptidase activity that has been useful for the acceleration of cheese ripening.  
Commercial proteases preparations containing carboxy-peptidase activity are known  
(e.g. FlavorPro 192 from Biocatalysts) but these always contain mixtures of amino-  
peptidase, endo-protease and possibly other protease activities.

25 Protease addition for cheese ripening can be done in various stages of cheese  
preparation. Preferably, the enzymes are added to the cheese milk prior to or together  
with the addition of the coagulant (e.g. chymosin). Addition at this point ensures a  
homogenous distribution of the enzymes throughout the cheese. Alternatively, the  
enzymes can be added in a later stage, e.g. during the salting stage in Cheddar making,  
30 but this introduces the risk of inhomogeneous enzyme distribution in the cheese and  
formation of so-called hot spots. For that reason, addition of the enzymes to the cheese  
milk is preferred. A disadvantage is that the majority of the enzyme (60-90%) is often not  
incorporated in the cheese curd, and is discarded in the whey fraction where it can give

rise to unwanted (proteolysis) that makes the whey less or not suited for further applications. Especially endo-proteases with significant activity at pH5-7 could cause such unwanted side-activities, but also amino peptidase that often have optimal activity in this pH range may give rise to formation of e.g. unwanted flavours. Another potential problem of especially endo-protease addition to the cheese milk is that they interfere with the coagulation process, giving rise to a-specific hydrolysis leading to reduction of cheese yield. Also amino-peptidases can cause yield-losses since they usually are well active at pH6-7, the usual pH range of cheese making. Proteases that are not or almost not active at pH values during cheese making, but which become active in the cheese are preferred because they will not interfere with the cheese making process and will not cause unwanted reactions in the whey.

#### **Description of the invention**

According to the present invention, accelerated cheese ripening can be obtained by the use of carboxy-peptidases. The carboxy-peptidase preparation should be free from endo-protease activity, and should be able to at least release amino acids that are important for cheese flavour formation, such as leucine, phenylalanine, valine and methionine. The carboxypeptidase is added at activity levels between 1 and 2500 CPG/g substrate (e.g. cheese milk), preferably 1-250 CPG/g substrate or more preferably 1-25 CPG/g substrate. CPG-units are defined in example 1. Protease activity is measured by the hydrolysis of a casein (6 g/L assay solution) at pH6.0, 4°C for 1 hour. 1 PU is the amount of enzyme that produces, in one minute, a (TCA-soluble) hydrolysate, which (280 nm) absorbance is equal to a tyrosin solution of 1 µM). The carboxypeptidase preparation is defined as free of endo-protease activity when the ratio of endo-protease activity (PU) / carboxypeptidase activity (CPG) in the preparation is less than 0.01, preferably less than 0.001 and most preferably less than 0.0005. The carboxy-peptidase is preferably a broad spectrum carboxy-peptidase that is able to release to majority of the natural amino acids from peptides or proteins. Broad spectrum carboxypeptidase is defined as an enzyme that is able to release at least 80% of the natural amino acids in amounts detectable by the method as described in example 3 of this application. Preferably the carboxypeptidase preparation contains a carboxy-peptidase in which 90%

of the carboxypeptidase activity is caused by a single enzyme, measured as described in example 1, but combinations of carboxypeptidases are also allowed.

We have surprisingly found that the use of a purified carboxy-peptidase CPD I (PEPG) from *Aspergillus niger* is well able to by its own accelerate cheese ripening. The enzyme has been described (Dal Degan, Ribadeau-dumas & Breddam, Appl. Environ. Microbiol (1992) 58, 2144-2152) and sequenced (Svendsen & Dal Degan, Bioch. Biophys. Acta (1998) 1387, 369-377). The carboxy-peptidase can also be used to accelerate cheese flavour development in EMCs. Other preferred applications of the carboxy-peptidase are in the field of flavour development in fermented foods like fermented sausages and beers in which, similar to the situation in cheese, availability of free amino acids like valine, leucine, isoleucine and phenylalanine as well as sulfur containing amino acids like methionine are known to be of particular importance.

#### **Example 1**

##### **Cloning of CPD-I (PEPG)**

The amino acid sequence of carboxypeptidase I (PEPG) is described (Svendsen & Dal Degan, Bioch. Biophys. Acta (1998) 1387, 369-377). Degenerate PCR primers were designed to clone the pepG gene from a genomic library from *Aspergillus niger* N400 (CBS 120.49), using methods known to the skilled person in the art. The gene was fused to the 3' end of the glucoamylase promotor. Analogous examples of fusions of structural genes to the glucoamylase promotor have been described (EP-A-0420358, EP-A-0463706 and WO99/38956). First the pepG structural gene was PCR amplified from a genomic fragment containing the gene and purified. Second, the promotor region of the glaA gene was PCR amplified using , at the 3' end, a primer that overlaps the 5' end of the pepG structural gene. Third, the two PCR fragments were fused via fusion PCR with an oligonucleotide primer 5' of the glaA promotor, and an oligonucleotide overlapping the stopcodon of pepG in the reverse direction. Fourth, the resulting fusion fragment was cloned in the *A niger* expression vector pGBTOP7 (WO99/38956), resulting in a fusion plasmid containing the glaA promotor, pepG structural gene and the glaA terminator. This plasmid was digested with HindIII and co-transformed with pGBBAAS-1 digested with Xho I to *Aspergillus niger* ISO502, essentially as described in WO 99/38956. Transformants selected for growth on acetamide plates were analysed using colony PCR to check for the presence of pepG expression cassette, using known

gene sequence was determined, and the genomic DNA sequence, coding sequence and corresponding amino acid sequence are given as SEQ ID NO.: 1, 2 and 3 respectively. *A. niger* pepG transformants were cultivated in shake flask using methods as described previously (WO 99/38956). After growth for 6 days at 34°C, supernatants were analysed on activity. Activity of PEPG was determined by adding 10 µl of the culture supernatant to a 990 µl of a solution containing 45 mM Na-acetate (pH4.5), 0.95 mM EDTA and 0.2 mM FA-Phe-Ala (obtained from Bachem). The change in optical density at 337 nm was followed. The decrease in optical density is a measure for the PEPG activity. One enzyme unit (1 CPG) is defined as the amount of enzyme needed to decrease the optical density at 337 nm by 1 absorbance unit per minute under the test conditions. The transformant showing the highest CPG value per ml was selected for PEPG expression.

### **Example 2**

#### **Purification of PEPG**

PEPG was purified from the culture broth of *Aspergillus niger* expressing the enzyme according to the method described (Dal Degan, Ribadeau-dumas & Breddam, Appl. Environ. Microbiol (1992) 58, 2144-2152) with the exception that the CABS-Sephadex step was omitted. The activity of the final substantially purified enzyme was established to be 150 CPG/ml, using the activity measurement as described in example 1.

### **Example 3**

#### **Determination of the substrate specificity of PEPG**

The substrate specificity of the purified PEPG was determined using substrates Z-Ala-X, in which Z is benzyloxycarbonyl and X is any of the amino acids (one letter code) A, D, E, F, G, H, I, K, L, M, N, P, Q, R, S, T, V, W, Y. All substrates were obtained from Bachem, except when X= Q or T, which substrates were obtained from PEPSCAN (The Netherlands) The enzyme specificity was determined at pH4.0 and 40 °C in solutions that contained 3 mM of the peptide substrates. The reaction was started by addition of the 5 µl enzyme solution ( 440 units/ml) to 95 µl of the reaction mixtures. Samples were taken for each substrate immediately at t=0 minutes and spotted on TLC-plates (Merck HPTLC [plates 20X10 Silica gel 60]), another sample was taken after 45

plates (Merck HPTLC [plates 20X10 Silica gel 60), another sample was taken after 45 minutes incubation and also spotted on the same TLC plate. As a control, the substrate solutions without the enzyme were spotted on the same TLC plate. The plate was stained for free amino groups by spraying with a ready to use ninhydrin spray (ACROS). Enzyme activity was rated from – (no activity), +/- (low activity) to + (little activity) to +++++ (very high activity). Very high activity (+++++) on a particular substrate is scored when all the substrate has already been converted at the t=0 sample. Results are as follows:

X	Activity score	Ref	X	Activity score	Ref
A	++	490	M	+++++	5820
C	nt	nt	N	+	nt
D	+	160	P	+/-	3
E	++	nt	Q	++	41
F	++++	nt	R	++	130
G	+	5	S	++	70
H	+	10	T	++	nt
I	++++	7090	V	+++++	3380
K	+++	200	W	+	nt
L	+++++	2950	Y	+++	nt

nt : not tested. Ref: data extracted from Dal Degan, Ribadeau-dumas & Breddam, Appl. Environ. Microbiol (1992) 58, 2144-2152. Numbers indicate kcat/kM values in min<sup>-1</sup> mM<sup>-1</sup>

The table shows that the cloned and purified PEPG is similar to the one described by Dal Degan in 1992, but there are some unexpected differences. The enzyme preferentially liberates the hydrophobic amino acids F, I, L, M and V. The preference of the cloned gene is, however, different from the one described by Dal Degan *et al*, which has highest preference for I whereas the cloned enzyme shows highest activity on L, M and V. Also, the purified enzyme is rather active on K, more active than e.g. on A and D, which is different from the data described by Dal Degan *et al*. Clearly, the carboxy-peptidase is has a very broad substrate specificity and is able to handle all amino acids except possibly C, which was not tested.

#### Example 4

##### Demonstration of accelerated cheese ripening of PEPG in mini cheese (Cheddar type).

Miniature cheeses were produced as described by Shakeel-Ur-Rehman et al. (Protocol for the manufacture of miniature cheeses in Lait, 78 (1998), 607-620). Raw cows milk was pasteurized by heating for 30 minutes at 63°C. The pasteurized milk was transferred to wide mouth plastic centrifuge bottles (200mL per bottle) and cooled to 31°C. Subsequently, 0.72 ml of starter culture DS 5LT1 (DSM Gist B.V., Delft, The Netherlands) was added to each of the 200 ml of pasteurised milk in the centrifuge bottles and the milk was ripened for 20 minutes. Then,  $\text{CaCl}_2$  (132  $\mu\text{L}$  of a 1  $\text{mol.L}^{-1}$  solution per 200mL ripened milk) was added, followed by addition of the coagulant (0.04 IMCU per ml). In case the experimental involved the use of PEPG, this enzyme was added together with the coagulant. The milk solutions were held for 40-50 minutes at 31°C until a coagulum was formed. The coagulum was cut manually by cutters of stretched wire, spaced 1 cm apart on a frame. Healing was allowed for 2 minutes followed by gently stirring for 10 minutes. After that, the temperature was increased gradually to 39°C over 30 minutes under continuous stirring of the curd / whey mixture. Upon reaching a pH of 6.2 the curd / whey mixtures were centrifuged at room temperature for 60 minutes at 1,700g. The whey was drained and the curds were held in a water bath at 36°C. The cheeses were inverted every 15 minutes until the pH had decreased to 5.2-5.3 and were then centrifuged at room temperature at 1,700g for 20 minutes. After manufacture the cheese were ripened at 12°C and sensory analysis was performed after 3 and 6 weeks of ripening by a minimum panel of 3 people. Several dosages of PEPG to the cheese milk were used: 0 (=control), 5, 50 and 500 CPG/200 ml cheese milk. The addition of PEPG clearly led to an overall increase in flavour intensity, leading to a more matured taste as compared to the control cheese. This was the case at all levels of PEPG addition, even though the lowest level of addition (5 CPG/200 ml) required 6 weeks ripening to give a clear effect. At the other doses (50 and 500 CPG/ml) the effect on taste was already obvious after 3 weeks of ripening. The results clearly showed that PEPG accelerates cheese flavour development, and that the effect was dose dependent, and that no off-tastes had developed.

### **Example 5**

#### **Demonstration of accelerated cheese ripening of PEPG in Gouda type cheese.**

Gouda cheeses were prepared in 200L vats using the starter culture DelvoTec® DX31D (obtained from DSM) and Maxiren600 (obtained from DSM; 55 IMCU/l milk;DSM) as the coagulant, using a standard Gouda cheese manufacture protocol known to the person skilled in the art. Raw milk was standardised to a casein to fat ratio of approximately 0.9 and pasteurised for 15 seconds at 72°C. PEPG was added immediately prior to addition of Maxiren at a level of 25 CPG/L. A control cheese was prepared from the same milk batch using the same manufacturing process but without addition of PEPG. After coagulation the curd was cut and stirred for 20 minutes at a low speed. Following this half the whey was replaced with warm water (40% of initial volume) to increase the temperature from 31°C to 36°C. After this the curd/whey mixture was stirred, at an increasing speed, for another 30-40 minutes until the curd was firm enough for drainage. The curd was hooped and left for half an hour before being placed into 5 kg moulds. The cheeses were pressed, using increasing pressure, for around 4 hours after which the cheeses were left to rest overnight. The following morning they were placed into brine for 24 hours. After allowing the cheeses to dry, a protective coating was placed on the cheese and the cheeses started their ripening period. During this period they were kept at 15°C, relative humidity of approximately 80%. The cheeses were regularly turned and coated during the ripening period. Cheeses were organoleptically assessed after 6 weeks and 3 months of ripening by using a trained panel consisting of a minimum number of 8 people. After 6 weeks, the cheese containing PEPG received significantly higher scores for sweetness as compared to the control cheese. After 3 months of ripening, the cheese containing PEPG was clearly different from the control cheese, showing significantly higher flavour (and also clearly different creaminess.. The cheese flavour of the cheese containing PEPG was pleasant and more mature, as compared to the control cheese. The experiment clearly showed that PEPG accelerates the cheese flavour development and that no off-tastes developed.

30. 01. 2004

(61)

CLAIMS

- 1 A process for the flavour development in a fermented food whereby a  
5 carboxypeptidase is used.
- 2 A process according to claim 1 whereby the fermented food is beer, sausage  
or cheese preferably cheese.
- 10 3 A process according to claim 1 or 2 whereby the carboxypeptidase activity is  
for at least 90% caused by a single enzyme.
- 4 A process according to any one of claims 1 to 3 whereby the ratio of  
15 endoprotease activity (PU) and carboxypeptidase activity (CPG) is less than  
0.01, preferably less than 0.001 and most preferably less than 0.0005
- 5 A process according to any one of claims 1 to 4, whereby the  
carboxypeptidase is CPD-1, preferably CPD-1 having the amino acid  
sequence of SEQ ID NO:3.





EPO - DG 1

30. 01. 2004

(61)

**ABSTRACT**

5

The present invention relates to a process for the flavour development in a fermented food whereby a carboxypeptidase is used.



30. 01. 2004

## SEQUENCE LISTING

(61)

<110> DSM IP Assets B.V.  
 <120> CARBOXYPEPTIDASE FOR CHEESE RIPENING  
 <130> 24178EP/P0/  
 <160> 3  
 <170> PatentIn version 3.1  
 <210> 1  
 <211> 1814  
 <212> DNA  
 <213> Aspergillus niger

<400> 1  
 atgcggggta ccacggcaat tgcttcatta ctactgggtcg gtcgggccac cagtctccaa 60  
 aatcctcatc gtcgggctgt tccgccccct ctctcgcatc gcagcgtagc gtctcgctcc 120  
 gtgcccgttg agcgccgaac caccgacttt gagtatttga ctaacaagac tgcaagtgcg 180  
 tgattccgtt ttttaactac cgcatttata gttctaagat caattgcagg attcctggtc 240  
 aatggcacia gcatccccga agtcgatttc gacgtcggcg agtcctacgc cggccttctc 300  
 cccaatacgc ccaactggcaa ttctagccta ttcttctggt tcttccccctc gcaaaatcca 360  
 gaggccagcg atgagggttag tggtcgctct gttttttccg gtcattgcgtc agccagctaa 420  
 caattaacaa agatcaccat ctgggtcaac ggcgggcccc gatgtagctc cctagacggc 480  
 ctgcttcaag agaacggccc attcctcttg cagcctggca cttacaagcc cgttcctaat 540  
 ccatactcat ggaccaacct caccaatgtg gtttacatcg accaaccgc cggcacaggc 600  
 ttctccccgg gccctcgac cgtaaataac gaggaagacg tggctgcca gttcaacagc 660  
 tggttcaagc acttcgtcga caccttcgac ctgcacggcc gcaaggtcta catcacgggt 720  
 gaaagctacg cgggcatgta cgtcccctac attgccgatg ccatgctgaa cgaggaggat 780  
 acaacctact tcaacttgaa gggtatccag atcaacgacc cgtccatcaa cagcgactcg 840  
 gtcattgatg actgtatgtt tcccttcata tacctccacc tccaccacca ccaccactaa 900  
 caacatcacc caccagcccc cgcggtccgc catctgaacc actacaacaa catcttccag 960  
 ctaaactcca ctttctctc ctacatcaac gccaaagccg acaagtgcgg ctacaacgcc 1020  
 ttctctgaca aagccatcac ctaccacccc ccagtcctt tccccaccgc cctgaaatc 1080  
 accgaagact gccaaagtctg ggacgaagtc gtcattggccg cctacgacat caaccctgc 1140  
 ttcaattact accacctgat cgacttctgc ccctacctct gggacgtgct cggcttcccc 1200

tccctcgctt	ccggcccaaa	caactacttc	aaccgctccg	acgtccagaa	gatcctgcac	1260
gtccctccaa	cggactactc	cgtgtgctcg	gagaccgtca	tcttcgogaa	cggcgacggc	1320
agcgacccca	gctcctgggg	tcccctaccc	agcgtcatcg	aacgcactaa	caacactatc	1380
atcggccacg	gctggctcga	ttacctcctc	ttcttgaacg	gctcgctcgc	cacaatccag	1440
aacatgacct	ggaacggtaa	gcaagggttc	cagcgtcctc	ccgtggaacc	gctcttcgtc	1500
ccttaccatt	atgggtctggc	tgagctgtac	tggggcgatg	agcctgaccc	gtataacctt	1560
gatgctggcg	ctggatacct	gggtacagcg	cataccgagc	gcggggttgac	tttcagctcg	1620
gtgtatattgt	ctggatcatg	taagtttatt	atatccccctt	ggaagcggta	tgatgaacgt	1680
tagagagtgc	tgactgttgc	ttcctcctcg	tgatagaaat	cccgcagtat	gttctctggcg	1740
cggcttaccg	ccagttggag	ttcctgctcg	gtaggattag	tagtctttcg	gcgaagggga	1800
actatacctc	ttga					1814

<210> 2

<211> 1563

<212> DNA

<213> *Aspergillus niger*

<400> 2

atgcgggta	ccacggcaat	tgcttcatta	ctactggctg	gctcggccac	cagtctccaa	60
aatcctcatc	gtcgggctgt	tccgcccctt	ctctcgcatc	gcagcgtagc	gtctcgctcc	120
gtgcccgttg	agcgccgaac	caccgacttt	gagtatttga	ctaacaagac	tgcaagattc	180
ctgggtcaatg	gcacaagcat	ccccgaagtc	gatttcgacg	tcggcgagtc	ctacgccggc	240
cttctcccca	atacgcccac	tggaatttct	agcctattct	tctggttctt	cccctcgcaa	300
aatccagagg	ccagcgatga	gatcaccatc	tggctcaacg	gcggccccgg	atgtagctcc	360
ctagacggcc	tgcttcaaga	gaacggccca	ttcctctggc	agcctggcac	ttacaagccc	420
gttcctaata	catactcatg	gaccaacctc	accaatgtgg	tttacatcga	ccaacccgcc	480
ggcacaggct	tctccccggg	cccctcgacc	gtaaataacg	aggaagacgt	ggctgcccag	540
ttcaacagct	ggttcaagca	cttcgtcgac	accttcgacc	tgacaggccg	caaggtctac	600
atcaccgggtg	aaagctacgc	gggcatgtac	gtcccctaca	ttgccgatgc	catgctgaac	660
gaggaggata	caacctactt	caacttgaag	ggtatccaga	tcaacgaccc	gtccatcaac	720
agcgactcgg	tcatgatgta	ctcccccgcc	gtccgccatc	tgaaccacta	caacaacatc	780
ttccagctaa	actccaattt	cctctctctac	atcaacgccca	aagccgacaa	gtgcgggtac	840

aacgccttcc togacaaagc catcacctac ccacccccca gtcccttccc caccgcccct 900  
 gaaatcaccg aagactgcc aagtctgggac gaagtcgtca tggccgccta cgacatcaac 960  
 ccctgcttca attactacca cctgatcgac ttctgcccct acctctggga cgtgctcggc 1020  
 ttcccctccc tcgcctccgg cccaaacaac tacttcaacc gctccgacgt ccagaagatc 1080  
 ctgcacgtcc ctccaacgga ctactccgtg tgctcggaga ccgtcatctt cgogaacggc 1140  
 gacggcagcg accccagctc ctgggggtccc ctaccagcg tcatcgaacg cactaacaac 1200  
 actatcatcg gccacggctg gctcgattac ctctcttct tgaacggctc gctcgccaca 1260  
 atccagaaca tgacctggaa cggtaagcaa gggttccagc gtcctcccgt ggaaccgctc 1320  
 ttcgctccctt accattatgg tctggctgag ctgtactggg gcgatgagcc tgaccggtat 1380  
 aaccttgatg ctggcgctgg atacctgggt acagcgcata ccgagcgcgg gttgactttc 1440  
 agctcgggtg atttgtctgg tcatgaaatc ccgcagtatg ttcttggtgc ggcttaccgc 1500  
 cagttggagt tctgctcgg taggattagt agtctttcgg cgaaggggaa ctatacctct 1560  
 tga 1563

<210> 3  
 <211> 520  
 <212> PRT  
 <213> *Aspergillus niger*

<400> 3

Met Arg Val Thr Thr Ala Ile Ala Ser Leu Leu Leu Val Gly Ser Ala  
 1 5 10 15

Thr Ser Leu Gln Asn Pro His Arg Arg Ala Val Pro Pro Pro Leu Ser  
 20 25 30

His Arg Ser Val Ala Ser Arg Ser Val Pro Val Glu Arg Arg Thr Thr  
 35 40 45

Asp Phe Glu Tyr Leu Thr Asn Lys Thr Ala Arg Phe Leu Val Asn Gly  
 50 55 60

Thr Ser Ile Pro Glu Val Asp Phe Asp Val Gly Glu Ser Tyr Ala Gly  
 65 70 75 80

Leu Leu Pro Asn Thr Pro Thr Gly Asn Ser Ser Leu Phe Phe Trp Phe  
 85 90 95

Phe Pro Ser Gln Asn Pro Glu Ala Ser Asp Glu Ile Thr Ile Trp Leu  
100 105 110

Asn Gly Gly Pro Gly Cys Ser Ser Leu Asp Gly Leu Leu Gln Glu Asn  
115 120 125

Gly Pro Phe Leu Trp Gln Pro Gly Thr Tyr Lys Pro Val Pro Asn Pro  
130 135 140

Tyr Ser Trp Thr Asn Leu Thr Asn Val Val Tyr Ile Asp Gln Pro Ala  
145 150 155 160

Gly Thr Gly Phe Ser Pro Gly Pro Ser Thr Val Asn Asn Glu Glu Asp  
165 170 175

Val Ala Ala Gln Phe Asn Ser Trp Phe Lys His Phe Val Asp Thr Phe  
180 185 190

Asp Leu His Gly Arg Lys Val Tyr Ile Thr Gly Glu Ser Tyr Ala Gly  
195 200 205

Met Tyr Val Pro Tyr Ile Ala Asp Ala Met Leu Asn Glu Glu Asp Thr  
210 215 220

Thr Tyr Phe Asn Leu Lys Gly Ile Gln Ile Asn Asp Pro Ser Ile Asn  
225 230 235 240

Ser Asp Ser Val Met Met Tyr Ser Pro Ala Val Arg His Leu Asn His  
245 250 255

Tyr Asn Asn Ile Phe Gln Leu Asn Ser Thr Phe Leu Ser Tyr Ile Asn  
260 265 270

Ala Lys Ala Asp Lys Cys Gly Tyr Asn Ala Phe Leu Asp Lys Ala Ile  
275 280 285

Thr Tyr Pro Pro Pro Ser Pro Phe Pro Thr Ala Pro Glu Ile Thr Glu  
290 295 300

Asp Cys Gln Val Trp Asp Glu Val Val Met Ala Ala Tyr Asp Ile Asn  
305 310 315 320

Pro Cys Phe Asn Tyr Tyr His Leu Ile Asp Phe Cys Pro Tyr Leu Trp  
325 330 335

Asp Val Leu Gly Phe Pro Ser Leu Ala Ser Gly Pro Asn Asn Tyr Phe  
340 345 350

Asn Arg Ser Asp Val Gln Lys Ile Leu His Val Pro Pro Thr Asp Tyr  
355 360 365

Ser Val Cys Ser Glu Thr Val Ile Phe Ala Asn Gly Asp Gly Ser Asp  
370 375 380

Pro Ser Ser Trp Gly Pro Leu Pro Ser Val Ile Glu Arg Thr Asn Asn  
385 390 395 400

Thr Ile Ile Gly His Gly Trp Leu Asp Tyr Leu Leu Phe Leu Asn Gly  
405 410 415

Ser Leu Ala Thr Ile Gln Asn Met Thr Trp Asn Gly Lys Gln Gly Phe  
420 425 430

Gln Arg Pro Pro Val Glu Pro Leu Phe Val Pro Tyr His Tyr Gly Leu  
435 440 445

Ala Glu Leu Tyr Trp Gly Asp Glu Pro Asp Pro Tyr Asn Leu Asp Ala  
450 455 460

Gly Ala Gly Tyr Leu Gly Thr Ala His Thr Glu Arg Gly Leu Thr Phe  
465 470 475 480

Ser Ser Val Tyr Leu Ser Gly His Glu Ile Pro Gln Tyr Val Pro Gly  
485 490 495

Ala Ala Tyr Arg Gln Leu Glu Phe Leu Leu Gly Arg Ile Ser Ser Leu  
500 505 510

Ser Ala Lys Gly Asn Tyr Thr Ser  
515 520



